

LOOKBACK TIME AND CHANDRA CONSTRAINS ON COSMOLOGICAL PARAMETERS

D. TONOIU, A. CARAMETE, L.A. POPA

Institute for Space Sciences, Atomistilor 409, PO Box MG-23, Magurele, RO 077125, Ilfov, Romania, E-mail: tonoiud@spacescience.ro, acaramete@spacescience.ro

Received July 29, 2010

Abstract. In this paper we combine the WMAP7 with lookback time and Chandra gas fraction data to constrain the main cosmological parameters and the equation of state for the dark energy. We find that the lookback time is a good measurement that can improve the determination of the equation of state for the dark energy with regard to other external data sets. We conclude that larger lookback time data set will further improve our determination of the cosmological parameters.

Key words: cosmological parameters, WMAP measurements, lookback time, Chandra gas fraction, Λ CDM cosmological model, dark energy equation of state, Monte Carlo simulations.

1. INTRODUCTION

The 7-year Wilkinson Microwave Anisotropy Probe (WMAP) data rigorously test the standard cosmological model placing constraints on its basic parameters. The WMAP measurements alone are not enough to break the degeneracy among some cosmological parameters or to place constraints on non-standard cosmological models. For example, measurements of cosmic microwave background (CMB) power spectrum alone do not strongly constrain the curvature of the universe characterized by the energy density parameter Ω_k . One needs to complement the CMB data with the luminosity or angular diameter distances measurements in order to constrain Ω_k because the astrophysical distances depend also on the expansion history of the universe [1].

There are conclusive evidences that the universe is in a state of accelerated expansion. The Hubble diagram of Type Ia supernovae (SNeIa, *e.g.* [2, 3, 4]), combined with CMB anisotropy measurements (*e.g.* [1, 5]), baryon acoustic oscillation (BAO) from the galaxy distribution data (*e.g.* [6, 7, 8, 9]), and galaxy cluster gas mass fraction measurements (*e.g.* [10, 11, 12]) support the idea that we live in a spatially-flat universe where nonrelativistic matter make almost 30% of the critical energy density while the rest is an unknown component called dark

energy, with negative effective pressure being responsible for the present phase of accelerated expansion of the universe [13].

Constraints on dark energy density parameter Ω_Λ and on its equation of state w (the ratio of pressure to energy density), can explain the nature of the repulsive force causing the acceleration of the universe. One possible explanation for this unknown component is an energy density constant in time and uniform in space. Such a cosmological constant (Λ) was originally postulated by Einstein to explain a static universe, later rejected when the expansion of the Universe was first detected and presently reinstated to account for the dark energy. Still, the computed value of Λ is expected to be 10^{120} larger than the observed one. Another cosmological scenario considers that dark energy is a dynamical scalar field with a time varying equation of state. An alternative explanation of the accelerating expansion of the Universe is that general relativity or the standard cosmological model is incorrect.

However, ground and space observations can not discriminate among different dark energy scenarios as the correct explanation of the observed accelerating universe: a cosmological constant, a dynamical scalar field or a modification of general relativity [14, 15].

The WMAP7-year data combined with other astrophysical measurements [1] place constraints on the dark energy. Assuming a flat universe ($\Omega_k \approx 0$), an accurate determination of the Hubble expansion rate (H_0) helps in improving the limit of the equation of state of the dark energy [16, 17]. In the paper of Komatsu *et al.* [1] from the joint analysis of WMAP7+BAO+H0 in the case of a time independent equation of state, a value of, $w = -1.10 \pm 0.14$ at 68% CL was obtained. Furthermore, adding high- z supernova data to their analysis a more stringent limit was obtained, $w = -0.98 \pm 0.053$ at 68% CL. However, this later result does not take into account the systematic errors in supernovae, which are comparable with the statistical errors [18, 19]. Also, combining the cluster abundance and 5-year WMAP data, Vikhlinin *et al.* [20] found that for a flat universe $w = -1.08 \pm 0.18$ at 68% CL. Furthermore, adding BAO [21] and the supernova data [22], they found $w = -0.991 \pm 0.09$ at 68% CL.

In this paper, we perform a joint analysis of the CMB-WMAP7 data, constraints on Hubble expansion rate inferred from the age of astrophysical objects using lookback time method (LBT) and measurements of the gas mass fraction of relaxed clusters from Chandra X-ray observatory (Chandra). Our aim is to investigate for several parameters of a given cosmological model which combination of this three data sets puts better constraints.

We choose the lookback time method because it has the advantage of using the ages of distant objects which are independent of each other so we can avoid biases present in techniques that use distances of primary or secondary indicators in the cosmic distance ladder method [13].

Moreover, we use Chandra measurements because they currently provide one of the best constraints on Ω_m and have the advantage of being remarkably simple and robust in terms of its underlying assumptions [10].

The paper is structured as follows: in section 2 we briefly present the lookback time method, in section 3 Chandra gas fraction experiment, in section 4 we describe the statistical analysis and present the data sets and finally, in the last sections, the results and conclusions.

2. LOOKBACK TIME METHOD

This time-based method uses ages of astrophysical objects (passively evolving galaxies or clusters of galaxies) which are independent of each other, to constrain cosmological parameters. The advantage of this method is that it avoids biases that are present in cosmic distance ladder method, offering an independent way to cross-check cosmological constraints obtained by using other methods [7].

The lookback time is defined as the difference between the present age of the Universe (t_0) and its age at redshift z , $t(z)$

$$t_L(z) = t_0 - t(z) = \int_0^z \frac{dz'}{(1+z')H(z)}, \quad (1)$$

where $H(z)$ is the Hubble parameter at redshift z .

We use the spatially-flat cosmological standard model Λ CDM, Λ denoting the fact that the dark energy component has a time independent unknown equation of state. In this model, the Hubble expansion rate as a function of redshift can be written as:

$$H(z) = H_0 \left[\Omega_m (1+z)^3 + (1-\Omega_m)(1+z)^{3(1+w)} \right]^{1/2}, \quad (2)$$

where Ω_m is the matter energy density parameter and w is the equation of state for the dark energy.

The observed lookback time $t_L^{obs}(z_i)$ (see the diagram presented in Fig. 1), to an object i at redshift z_i is defined as:

$$t_L^{obs}(z_i) = t_0^{obs} - t_i(z_i) - df, \quad (3)$$

where t_0^{obs} is the measured age of the Universe and $t_i(z_i)$ is the age of the object (a passively evolving galaxy or a cluster) defined as the difference between the age of the universe at redshift z_i and the age of the universe at the redshift of object formation, z_f :

$$t_i(z_i) = t(z_i) - t(z_f) = t_L(z_f) - t_L(z_i) = \int_{z_i}^{z_f} \frac{dz'}{(1+z')H(z')}. \quad (4)$$

The third term in equation (3), df is a delay factor that encodes our ignorance regarding the formation redshift of the object z_f and is defined as:

$$df = t_0^{obs} - t_L(z_f).$$




Fig. 1 – The time evolution of the lookback time and the age of the universe as a function of redshift [17] (see also the text).

The lookback time method has the important feature that the age of distant objects are independent of each other and we can avoid biases present in techniques that use distances of primary or secondary indicators in the cosmic distance ladder method [13].

3. CHANDRA X-RAY CLUSTER GAS FRACTION

It was shown that the gas fraction (f_{gas}) in X-ray luminous, dynamically relaxed clusters measured with the Chandra X-ray Observatory, helps in better constraining the cosmological parameters [10, 23]. Following [24, 25], the matter content of the largest clusters of galaxies provides an almost fair sample of the total matter of the universe. Measurements of the ratio of baryonic and total mass in clusters of galaxies should closely match the ratio of Ω_b and Ω_m . The matter

energy density parameter Ω_m can be determined by combining measurements of the baryonic mass fraction in the largest galaxy clusters with accurate determinations of $\Omega_b h^2$ from cosmic nucleosynthesis and CMB experiments, and a reliable measurement of the Hubble constant, H_0 .

The X-ray clusters gas fraction model fitted to a reference Λ CDM cosmology is [10]

$$f_{gas}^{\Lambda CDM}(z) = \frac{KA\gamma b(z)}{1+s(z)} \left(\frac{\Omega_b}{\Omega_m} \right) \left[\frac{d_A^{\Lambda CDM}(z)}{d_A(z)} \right]^{1.5}, \quad (5)$$

where $d_A(z)$ and $d_A^{\Lambda CDM}(z)$ are the angular diameter distances to the clusters in the current test model XCDM and the reference standard cosmological model Λ CDM.

$$d_A = \frac{c}{H_0(1+z)\sqrt{\Omega_k}} \sinh \left(\sqrt{\Omega_k} \int_0^z \frac{H_0}{H(z)} dz \right). \quad (6)$$

The systematic uncertainties in the Chandra experiment are parameterized by the angular correction A, the non-thermal pressure support in the clusters γ , the baryonic mass fraction in stars $s(z) = s_0(1 + \alpha_s z)$, the bias factor $b(z) = b_0(1 + \alpha_b z)$ and an overall calibration parameter K for the residual uncertainty in the accuracy of the instrument calibration and X-ray modeling.

4. ANALYSIS

We perform our analysis in the framework of the extended XCDM cosmological model, described by 7 + 1 free parameters:

$$\Theta = (\underbrace{\Omega_b h^2, \Omega_{DM} h^2, \theta_s, \tau, w, n_s, A_s}_{standard}, df), \quad (7)$$

assuming uniform priors for all the parameters.

Here $\Omega_b h^2$ and $\Omega_{DM} h^2$ are the baryonic and dark matter energy density parameters, θ_s is the ratio of the sound horizon distance to the angular diameter distance, τ is the reionization optical depth, $w = \frac{p}{\rho}$ is the equation of state parameter (p and ρ are the pressure and energy density of the dark energy), n_s is the scalar spectral index of the primordial density perturbation power spectrum, A_s is its amplitude at the pivot scale and df is the lookback time delay factor. Table 1 presents the parameters of our model, their fiducial values and the prior ranges adopted in the analysis.

Table 1

The parameters of our model, their fiducial values and the prior ranges adopted in the analysis

Parameter	Fiducial value	Prior range
$\Omega_b h^2$	0.0223	0.005 \rightarrow 0.1
$\Omega_{DM} h^2$	0.105	0.01 \rightarrow 0.99
θ_s	1.04	0.5 \rightarrow 10
τ	0.09	0.01 \rightarrow 0.8
w	-1	-2 \rightarrow 0
n_s	0.95	0.5 \rightarrow 1.5
$\ln[10^{10} A_s]$	3	2.7 \rightarrow 4
df [Gyr]	1.5	0 \rightarrow 3

We modified the CosmoMC Monte Carlo Markov Chain (MCMC) public package [26] for our extended θ parameter space to sample from the posterior distribution giving the following experimental datasets:

The **WMAP7** temperature and polarization CMB latest measurements [1].

The **Lookback time (LBT)** measurements. From Ref. [27] we use the ages of six galaxy clusters in the redshift range $0.10 \leq z \leq 1.27$. The standard deviation uncertainty for this age measurements is 1 Gyr. We also use the age of 32 passively evolving galaxies from Ref. [28] in the redshift interval $0.117 \leq z \leq 1.845$. The error for this sample (1σ) is 12% of the age measurements. Therefore, we have 38 measurements of $t_i(z_i)$ with uncorrelated uncertainties σ_i . For the age of the universe we use the WMAP estimate, $t_0^{obs} = (13.69 \pm 0.13)$ Gyr [5].

Then, we compute the χ^2 function, for each cosmological model described by the set of parameters given in equation (7) as:

$$\chi_{LBT}^2(\theta) = \sum_{i=1}^{38} \frac{(t_L(\theta) - t_L^{obs}(z_i, df))^2}{\sigma_i^2 + \sigma_{t_0^{obs}}^2} + \frac{(t_0(\theta) - t_0^{obs})^2}{\sigma_{t_0^{obs}}^2}, \quad (8)$$

where t_L and t_0 are the theoretical predicted values of the lookback time and of the age of the universe and t_L^{obs} and t_0^{obs} are the corresponding measured values. Also, σ_i is referring to the one standard error of the experimental data and $\sigma_{t_0^{obs}}$ is the uncertainty in the estimate of t_0 .

Chandra X-ray gas fraction. We used fgas data following the work from Ref. [10, 23, 29]. We modified the CosmoMC Monte Carlo Markov Chain public package [26] to allow the use of fgas data when exploring the cosmological parameter space, implementing fgas module presented in Ref. [30].

Our implementation of this new module into CosmoMC package considers all seven parameters related to systematic uncertainties as being constant and having the following values: $A = 0.2$, $\gamma = 1.05$, $s_0 = 0.16$, $\alpha_s = 0$, $b(z) = 0.824$, $\alpha_b = 0$ and $K = 1$. This is different from the approach described in [10], where either Gaussian or linear uncertainties are taken into consideration and after that, the seven parameters are added to the original list of 13 possible free parameters. In this manner we manage to decrease the computational costs, especially when trying to constrain non-standard models using also the Chandra data.

The χ^2 function we used for f_{gas} data has the form:

$$\chi_{CH}^2 = \sum_{i=1}^{42} \frac{(f_{gas}^{\Lambda CDM}(z) - f_{gas}^{\text{exp}}(z))^2}{\sigma_{f_{gas}}^{\text{exp}}(z)^2} + \frac{(\eta - 0.214)^2}{(0.022)^2}, \quad (9)$$

where, $f_{gas}^{\Lambda CDM}$ is the gas fraction fitted to a reference ΛCDM cosmological model and is given by (5) and $f_{gas}^{\text{exp}}(z)$, $\sigma_{f_{gas}}^{\text{exp}}(z)$ are the experimental values of gas fraction and the associated errors. In the above equation, η is the slope of the $f_{gas}(z)$ in a region with a radius for which the mean enclosed mass density is 2 500 times the critical density of the universe at the redshift of the cluster, as measured for the reference ΛCDM standard cosmological model [10]. The difference between the experimental slope η and its reference value is normalized to its expected standard deviation squared.

WMAP7, LBT and CHANDRA data sets (WMAP7+LBT+CHANDRA) are combined by multiplying the likelihoods.

5. RESULTS

In order to see, for several parameters of the chosen cosmological model ΛCDM , which data provides better constrains we run the modified CosmoMC package on a parallel computing system by using 64 independent chains, for the following combinations of data sets: WMAP7, WMAP7+LBT, WMAP7+CHANDRA and WMAP7+LBT+CHANDRA. We impose for each case the Gelman & Rubin convergence criterion [31].

We present in Table 2, the mean value and 1σ error at a 68% CL, obtained from our analysis, for the following cosmological parameters: $\Omega_b h^2$, $\Omega_{DM} h^2$, Ω_Λ , Ω_m , n_s , w , H_0 , Age(Gyr), $\log[10^{10} A_s]$. In the first column, we present the results for the reference WMAP-WCDM model¹ and in the second column, our simulation with WMAP7 data alone. Then, in the next two columns we present the results

¹ http://lambda.gsfc.nasa.gov/product/map/dr4/params/wcdm_sz_lens_wmap7.cfm

obtain from the combination of WMAP7 with LBT and Chandra data and finally, in the last column, we show the results when all three data sets are joined. Comparing with the WMAP-WCDM reference model, we conclude that our computation best constrain the $\Omega_{DM} h^2$, Ω_Λ , Ω_m parameters as expected, if we use all three data sets (WMAP7, LBT, Chandra) and the parameters $\Omega_b h^2$, n_s , w , H_0 , Age, $\log[10^{10} A_s]$ are better constrained with WMAP7 and Chandra data.

In Fig. 2 we show the correlation between some cosmological parameters, $(\Omega_m, \Omega_\Lambda)$, (Ω_m, w) , (H_0, Ω_m) , (H_0, w) , (Age, w) , (Ω_Λ, w) . This are the joint two-dimensional marginalized distributions with 68%, 95%, 99% CL for the following combinations of data sets: WMAP7 data only, WMAP7+LBT, WMAP7+Chandra and WMAP7+LBT+Chandra. The plane $\Omega_m - \Omega_\Lambda$ is better constrained when we combine all three data sets. Also for others correlations, WMAP7 and Chandra combination is the optimal one. Although adding LBT data set to the analysis we obtain better constraints upon all cosmological parameters, the central values of the distributions are shifted when comparing with WMAP-WCDM reference model, so we prefer to use only the WMAP7+Chandra data sets when we want to see the correlation between the equation of state of dark energy and other cosmological parameters: Ω_m , H_0 , Age(Gyr), Ω_Λ and also, the correlation between H_0 and Ω_m .

We have shown that, for some parameters of a cosmological model with an unknown time independent equation of state, adding LBT to Chandra and WMAP7 data, leads to an improvement of their error bars. In the future, we expect that a larger LBT data set will further improve constrains on cosmological parameters.

Table 2

The mean and 1σ uncertainty for cosmological parameters in WMAP7, LBT, and Chandra combined data analysis

	WCDM- WMAP7	W7	W7+LBT	W7+Chandra	W7+LBT+ Chandra
$\Omega_b h^2$	$0.02258^{+0.00063}_{-0.00062}$	$0.02259^{+0.00058}_{-0.00058}$	$0.02137^{+0.00073}_{-0.00068}$	$0.02202^{+0.00047}_{-0.00047}$	$0.01909^{+0.00046}_{-0.00046}$
$\Omega_{DM} h^2$	$0.1112^{+0.0058}_{-0.0058}$	$0.1111^{+0.0055}_{-0.0055}$	$0.0766^{+0.0039}_{-0.0044}$	$0.1192^{+0.0038}_{-0.0038}$	$0.0945^{+0.0024}_{-0.0025}$
Ω_Λ	$0.741^{+0.095}_{-0.099}$	$0.711^{+0.024}_{-0.025}$	$0.763^{+0.010}_{-0.010}$	$0.694^{+0.016}_{-0.016}$	$0.725^{+0.007}_{-0.007}$
Ω_m	$0.259^{+0.099}_{-0.095}$	$0.289^{+0.025}_{-0.024}$	$0.237^{+0.010}_{-0.010}$	$0.306^{+0.016}_{-0.016}$	$0.275^{+0.007}_{-0.007}$
n_s	$0.964^{+0.015}_{-0.015}$	$0.967^{+0.014}_{-0.014}$	$0.973^{+0.018}_{-0.016}$	$0.951^{+0.011}_{-0.011}$	$0.909^{+0.010}_{-0.010}$
w	$-1.12^{+0.42}_{-0.43}$	$-0.92^{+0.10}_{-0.10}$	$-0.65^{+0.04}_{-0.03}$	$-1.04^{+0.11}_{-0.11}$	$-0.85^{+0.02}_{-0.02}$
$H_0[\text{km} \cdot \text{s}^{-1} \text{Mpc}^{-1}]$	75^{+15}_{-14}	$68^{+3.9}_{-2.7}$	$64^{+0.2}_{-0.3}$	$68^{+2.2}_{-2.1}$	$64^{+0.2}_{-0.2}$
Age [Gyr]	$13.75^{+0.29}_{-0.27}$	$13.81^{+0.11}_{-0.11}$	$14.34^{+0.10}_{-0.10}$	$13.88^{+0.1}_{-0.1}$	$14.60^{+0.09}_{-0.09}$
$\log[10^{10} A_s]$	-	$3.078^{+0.035}_{-0.035}$	$2.945^{+0.040}_{-0.040}$	$3.086^{+0.032}_{-0.033}$	$2.942^{+0.032}_{-0.032}$




Fig. 2 – Joint 2D marginalized distributions at 68%, 95%, 99% CL showing the correlation between $(\Omega_m, \Omega_\Lambda)$, (Ω_m, w) , (H_0, Ω_m) , (Age, w) , (Ω_Λ, w) . The blue contour is for WMAP only data set, the red contour for WMAP+LBT, the yellow for WMAP7+Chandra and the green contour for WMAP7+LBT+Chandra data sets.

6. CONCLUSIONS

The aim of our paper was to analyze how the LBT and Chandra data sets combined with WMAP7 measurements can improve the determination of the equation of state for the dark energy and also of the other parameters of the standard cosmological model. We choose the LBT external data set because it contains information independent of each other and we can avoid biases present in the cosmic distance ladder method and the Chandra data set because it provides one of the best constrains on the matter density parameter Ω_m .

Our contribution consists in implementing LBT and Chandra new modules into the public Monte Carlo Markov Chain package. We run the modified CosmoMC package for the following combinations of data sets: WMAP7, WMAP7+LBT, WMAP7+Chandra, WMAP7+LBT+Chandra. We found that, the physical baryonic density parameter, the scalar spectral index, the equation of state

for the dark energy, the Hubble expansion rate, the age of the universe and the amplitude at the pivot scale are best constrain using WMAP7+Chandra data sets. All the other considered parameters, the physical dark matter, dark energy and matter density parameters and the plane $\Omega_m - \Omega_\Lambda$ are best constrain when we combine WMAP7+LBT+Chandra.

Moreover, we conclude that the looking back time is trustful measurement and in the future a larger LBT data set can further improve the determination of parameters we investigate.

Acknowledgements. This work is supported by the ESA/PECS Contract “Scientific exploitation of Planck-LFI data” and by CNCSIS Contract 539/2009.

REFERENCES

1. E. Komatsu, et al., *Seven-Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Cosmological Interpretation*, ApJS, **180**, 330 (2009).
2. M. Hicken, et al., *Improved Dark Energy Constraints from ~100 New CfA Supernova Type Ia Light Curves*, ApJ, **700**, 1097–1140 (2009).
3. A. Shafieloo, V. Sahni, A.A. Starobinsky, *Is cosmic acceleration slowing down?*, Phys. Rev. D, **80**, 101301 (2009).
4. A.C.C. Guimaraes, J.V. Cunha, J.A.S. Lima, *Bayesian Analysis and Constraints on Kinematic Models from Union SNIa*, JCAP, **0910**, 010 (2009).
5. J. Dunkley, et al., ApJS, **180**, 306 (2009).
6. W.J. Percival, et al., MNRAS, **381**, 1053 (2007).
7. L. Samushia, B. Ratra, *Constraints on dark energy from baryon acoustic peak and galaxy cluster gas mass measurements*, ApJS, **703**, 1904-1910 (2009).
8. E. Gasztañaga, A. Cabré, L. Hui, *Clustering of Luminous Red Galaxies IV: Baryon Acoustic Peak in the Line-of-Sight Direction and a Direct Measurement of $H(z)$* ; arXiv:0807.3551v5 [astro-ph], 2009.
9. Y. Wang, *Clarifying Forecasts of Dark Energy Constraints from Baryon Acoustic Oscillations*; arXiv:0904.2218 [astro-ph.CO], 2009.
10. S. W. Allen, et al., *Improved constraints on dark energy from Chandra X-ray observations of the largest relaxed galaxy clusters*, MNRAS, **383**, 879 (2008).
11. L. Samushia, B. Ratra, *Constraints on Dark Energy from Galaxy Cluster Gas Mass Fraction Versus Redshift Data*, ApJ, **680**, L1 (2008).
12. S. Ettori, et al., *The cluster gas mass fraction as a cosmological probe: a revised study*; arXiv:0904.2740 [astro-ph.CO], 2009.
13. L. Samushia et al., *Constraints on dark energy from the lookback time versus redshift test*; arXiv:0906.2734v1 [astro-ph.CO], 2009.
14. A. Albrecht, et al., *Report of the Dark Energy Task Force*; arXiv:astro-ph/0609591v1, 2006.
15. P.J.E. Peebles, B. Ratra, *The Cosmological Constant and Dark Energy*, arXiv:astro-ph/0207347v2, 2002.
16. D. N. Spergel et al., *First Year Wilkinson Microwave Anisotropy Probe (WMAP) Observations: Determination of Cosmological Parameters*, ApJ, **148**, 175 (2003).
17. W. Hu, *Dark Energy Probes in Light of the CMB*, Astronomical Society of the Pacific Conference Series, Vol. 339, 2005.
18. R. Kessler et al., *First-year Sloan Digital Sky Survey-II (SDSS-II) Supernova Results: Hubble Diagram and Cosmological Parameters*, ApJS, **185**, 32–84 (2009).

19. M. Hicken et al., *Improved Dark Energy Constraints from ~100 New CfA Supernova Type Ia Light Curves*, ApJ, **700**, 1097 (2009).
20. A. Vikhlinin et al., *Chandra Cluster Cosmology Project III: Cosmological Parameter Constraints*, ApJ, **692**, 1060 (2009).
21. D.J. Eisenstein et al., *Detection of the Baryon Acoustic Peak in the Large-Scale Correlation Function of SDSS Luminous Red Galaxies*, ApJ, **633**, 560 (2005).
22. T.M. Davis et al., *Scrutinizing Exotic Cosmological Models Using ESSENCE Supernova Data Combined with Other Cosmological Probes*, ApJ, **666**, 716 (2007).
23. S.W. Allen et al., *Constraints on dark energy from Chandra observations of the largest relaxed galaxy clusters*, MNRAS, **353**, 457A; astro-ph/0405340, 2004.
24. S.D.M. White et al., *Nature, The Baryon Content of Galaxy Clusters: A Challenge to Cosmological Orthodoxy*, **366**, 429 (1993).
25. V.R. Eke et al., *The Evolution of X-Ray Clusters in a Low-Density Universe*, ApJ, **503**, 569 (1998).
26. A. Lewis, S. Bridle, *Cosmological parameters from CMB and other data: a Monte-Carlo approach*, Phys. Rev. D., **66**, 103511 (2002).
27. S. Capozziello, et al., *Constraining dark energy models using the lookback time to galaxy clusters and the age of the universe*, Phys. Rev. D, **70**, 123501 (2004).
28. J. Simon, L. Verde, R. Jimenez, *Constraints on the redshift dependence of the dark energy potential*, Phys. Rev. D, **71**, 123001 (2005).
29. D. Rapetti, S.W. Allen, J. Weller, *Constraining Dark Energy with X-ray Galaxy Clusters, Supernovae and the Cosmic Microwave Background*, MNRAS, **360**, 55R (2005).
30. Rapetti web page: <http://www.stanford.edu/drapetti/fgas> module
31. A. Gelman, D. Rubin, *Inference from iterative simulation using multiple sequences*, Statistical Science, **7**, 457 (1992).