

# NEW RHEOLOGICAL MODELS WITHIN LOCAL FRACTIONAL DERIVATIVE

XIAO-JUN YANG<sup>1,2</sup>, FENG GAO<sup>1,2</sup>, H. M. SRIVASTAVA<sup>3,4</sup>

<sup>1</sup>School of Mechanics and Civil Engineering, China University of Mining and Technology,  
Xuzhou 221116, Peoples Republic of China  
Email: *dYangxiaojun@163.com*

<sup>2</sup>State Key Laboratory for Geomechanics and Deep Underground Engineering, China University of  
Mining and Technology,  
Xuzhou 221116, Peoples Republic of China  
Email: *jsppw@sohu.com*

<sup>3</sup>Department of Mathematics and Statistics, University of Victoria,  
Victoria, British Columbia V8W3R4, Canada  
Email: *harimsri@math.uvic.ca*

<sup>4</sup>Department of Medical Research, China Medical University Hospital,  
Taichung 40402, Taiwan, Republic of China

*Compiled April 21, 2017*

In this article, two local fractional rheological models via spring and dashpot elements in the one-dimensional case are proposed for the first time. The creep and relaxation behaviors of the fractal Maxwell fluid and Kelvin-Voigt solid are discussed. The constitutive relationships of the fractal mechanical elements are formulated by using local fractional calculus. The viscoelastic characteristics of the real materials are efficiently demonstrated and illustrated with the fractal charts.

*Key words:* rheology, fractal Maxwell fluid, fractal Kelvin-Voigt solid, local fractional derivative.

*PACS:* 87.19.rh, 61.43.Hv

## 1. INTRODUCTION

Fractional calculus (FC) has been successfully used in recent years in order to model the mathematical problems in real materials (see, for details, Refs. [1–12] and the works cited therein). For example, the representation of the constitutive relationship (RCR) of the fractional mechanical element with fractional derivative (FD) of the Riemann-Liouville type was given in [2]; its generalized version was proposed in [13]. The RCR with FC of the Liouville-Caputo type was considered in [14,15]. The RCR via fractional derivative of the Caputo-Fabrizio type was presented in [16]. The historical perspective in the fractional linear viscoelasticity was reported for example in Refs. [17–21].

Recently, local fractional calculus (LFC) has played important roles in the de-

scription of the fractal diffusion [22], fluid flow [23], shallow water surfaces [24], free damped vibrations [25] and other areas (see, for example, [26–35] and related references cited therein). To date, LFC has not been utilized to consider the representations of the rheological behaviors in real materials. Motivated by the previous idea, from mathematical viewpoint, the chief objective of the present paper is to propose the local fractional rheological models (LFRMs) for real materials via fractal spring and dashpot elements involving local fractional derivative (LFD) and to investigate the creep and relaxation behaviors of the fractal Maxwell fluid and Kelvin-Voigt solid with the aid of the local fractional Laplace transform (LFLT).

This article is structured as follows. In Sec. 2, the LFD and LFLT of the non-differentiable function are formulated. In Sec. 3, the fractal spring and dashpot elements within LFD are proposed. In Sec. 4, the fractal Maxwell element via LFD is analyzed. The Kelvin-Voigt element via LFD is reported in Sec. 5. Finally, in Sec. 6, the conclusion is outlined briefly.

## 2. MATHEMATICAL FUNDAMENTALS

In this Section, the concepts of the LFD and LFLT of the non-differentiable functions, which are utilized in this paper, are presented.

The LFD of the non-differentiable function  $\Lambda_\vartheta(t)$  of fractal order  $\vartheta$  ( $0 < \vartheta < 1$ ) at the point  $t = t_0$  is defined as follows (see, for details, [22–27, 31–35]):

$$\frac{\partial^\vartheta \Lambda_\vartheta(t_0)}{\partial t^\vartheta} = \lim_{t \rightarrow t_0} \frac{\Delta^\vartheta(\Lambda_\vartheta(t) - \Lambda_\vartheta(t_0))}{(t - t_0)^\vartheta}, \quad (1)$$

where

$$\Delta^\vartheta(\Lambda_\vartheta(t) - \Lambda_\vartheta(t_0)) \cong \Gamma(1 + \vartheta) \Delta[\Lambda_\vartheta(t) - \Lambda_\vartheta(t_0)].$$

The LFDs of the non-differentiable functions (see, for example, [22, 25, 26]) are listed in Table 1 below.

Table 1.

The LFDs of the non-differentiable functions.

Non-differentiable functions	LFDs
$t^{k\vartheta}/\Gamma(1 + k\vartheta)$ ( $k \in \mathbb{N}$ ) $M_\vartheta(\kappa t^\vartheta)$	$t^{(k-1)\vartheta}/\Gamma[1 + (k-1)\vartheta]$ $\kappa M_\vartheta(\kappa t^\vartheta)$

The LFLT of the non-differentiable function  $\psi_\vartheta(t)$  is given by [26]:

$$\psi_\vartheta(s) = {}_0I_\infty^{(\vartheta)} \left[ M_\vartheta \left( -s^\vartheta t^\vartheta \right) \psi_\vartheta(t) \right] \quad (0 < \vartheta \leq 1), \quad (2)$$

where  ${}_0I_\infty^{(\vartheta)}$  is the local fractional integral operator (see, for details, [24, 26]) and  $s$  is

the parameter of the local fractional Laplace integral operator (see [26]).

The LFLTs of the non-differentiable functions (see, e.g., [26]) are listed in Table 2.

Table 2.

The LFLTs of the non-differentiable functions.

Non-differentiable functions	LFLTs
$t^{k\vartheta}/\Gamma(1+k\vartheta)$ ( $k \in \mathbb{N}$ )	$1/s^{(k+1)\vartheta}$
1	$1/s^\vartheta$
$M_\vartheta(\kappa t^\vartheta)$	$1/(s^\vartheta - \kappa)$

### 3. FRACTAL SPRING AND DASHPOT ELEMENTS OF THE NON-DIFFERENTIABILITY TYPE

The representations of the fractal mechanical elements including the fractal spring and dashpot of the non-differentiability type are given in this Section.

As a matter of fact, the fractal spring of the non-differentiability type is the fractal elastic (or fractal storage) element illustrated in Fig. 1. This linear model, when represented as a perfect fractal elastic solid, obeys the Hooke's law of the non-differentiability type:

$$\sigma_\vartheta(t) = E_\vartheta \varepsilon_\vartheta(t), \quad (3)$$

where  $E_\vartheta$  is the Young's modulus of the fractal material,  $\varepsilon_\vartheta(t)$  is the fractal strain, and  $\sigma_\vartheta(t)$  is the fractal stress.




Fig. 1 – The representation of the fractal spring element.

The fractal dashpot within LFD is the fractal viscous (or fractal dissipative) element depicted in Fig. 2. The constitutive equation of non-differentiability type used to describe the viscous flow, which is expressed as the Newton's law of non-differentiability type, is given by:

$$\sigma_\vartheta(t) = \eta_\vartheta \frac{d^\vartheta \varepsilon_\vartheta(t)}{dt^\vartheta}, \quad (4)$$

where  $\eta_\vartheta$  is the fractal viscosity,  $\varepsilon_\vartheta(t)$  is the fractal strain, and  $\sigma_\vartheta(t)$  is the fractal stress.




Fig. 2 – The representation of the fractal dashpot element within LFD.

With the aids of the Boltzmann superposition principle and causal histories for  $t \in (0, \infty)$ , the creep and the relaxation representations of the non-differentiability type are expressed by

$$\varepsilon_\vartheta(t) = \sigma_\vartheta(0^+) J_\vartheta(t) + \frac{1}{\Gamma(1+\vartheta)} \int_0^t J_\vartheta(t-\tau) \frac{d^\vartheta \sigma_\vartheta(\tau)}{d\tau^\vartheta} (d\tau)^\vartheta \quad (5)$$

and

$$\sigma_\vartheta(t) = \varepsilon_\vartheta(0^+) G_\vartheta(t) + \frac{1}{\Gamma(1+\vartheta)} \int_0^t G_\vartheta(t-\tau) \frac{d^\vartheta \varepsilon_\vartheta(\tau)}{d\tau^\vartheta} (d\tau)^\vartheta, \quad (6)$$

respectively

From Eq. (5), the fractal creep compliance of the non-differentiability type is defined as follows:

$$J_\vartheta(t) = \frac{\varepsilon_\vartheta(t)}{\sigma_0}, \quad (7)$$

where  $\sigma_\vartheta(0^+) = \sigma_0$  is the initial stress condition.

From Eq. (6), the fractal relaxation modulus of the non-differentiability type is defined by:

$$G_\vartheta(t) = \frac{\sigma_\vartheta(t)}{\varepsilon_0}, \quad (8)$$

where  $\varepsilon_\vartheta(0^+) = \varepsilon_0$  is the initial strain condition.

Applying the LFLT to Eq. (5) and Eq. (6), we have

$$\varepsilon_\vartheta(s) = \frac{1}{s^\vartheta} J_\vartheta(s) \sigma_\vartheta(s) \quad (9)$$

and

$$\sigma_\vartheta(s) = \frac{1}{s^\vartheta} G_\vartheta(s) \varepsilon_\vartheta(s), \quad (10)$$

which lead us to

$$G_\vartheta(s) J_\vartheta(s) = \frac{1}{s^{2\vartheta}}, \quad (11)$$

where  $\sigma_\vartheta(s)$ ,  $\varepsilon_\vartheta(s)$ ,  $G_\vartheta(s)$  and  $J_\vartheta(s)$  are the LFLT's of  $\sigma_\vartheta(t)$ ,  $\varepsilon_\vartheta(t)$ ,  $G_\vartheta(t)$ , and  $J_\vartheta(t)$ , respectively.

From Eq. (11), the relationship between the fractal creep and the relaxation functions is of the following form:

$$\frac{1}{\Gamma(1+\vartheta)} \int_0^t G_\vartheta(t-\tau) J_\vartheta(\tau) (d\tau)^\vartheta = \frac{t^\vartheta}{\Gamma(1+\vartheta)} \quad (12)$$

or

$$\frac{1}{\Gamma(1+\vartheta)} \int_0^t J_\vartheta(t-\tau) G_\vartheta(\tau) (d\tau)^\vartheta = \frac{t^\vartheta}{\Gamma(1+\vartheta)}. \quad (13)$$

#### 4. FRACTAL MAXWELL ELEMENT VIA LFD

The fractal Maxwell element, when adopted to represent the fractal fluid, is made up of the fractal spring and dashpot in series as illustrated in Fig. 3.




Fig. 3 – The representation of the fractal Maxwell fluid involving LFD.

The constitutive equation of the fractal Maxwell model within LFD can be written in the form:

$$\frac{d^\vartheta \varepsilon_\vartheta(t)}{dt^\vartheta} = \frac{\sigma_\vartheta(t)}{\eta_\vartheta} + \frac{1}{E_\vartheta} \frac{d^\vartheta \sigma_\vartheta(t)}{dt^\vartheta}, \quad (14)$$

where  $\eta_\vartheta$  is the fractal viscosity,  $E_\vartheta$  is the Young's modulus,  $\varepsilon_\vartheta(t)$  is the fractal strain, and  $\sigma_\vartheta(t)$  is the fractal stress.

When the fractal body is subjected to the following condition:

$$\sigma_\vartheta(t) = \sigma_0 H(t), \quad (15)$$

where  $H(t)$  is the unit step function defined on fractal sets (see [26]) and  $\sigma_0$  is the a constant stress, Eq. (14) becomes

$$\frac{d^\vartheta \varepsilon_\vartheta(t)}{dt^\vartheta} = \frac{\sigma_0(t) H(t)}{\eta_\vartheta}. \quad (16)$$

Upon taking the LFLT of both sides of Eq. (16), we have

$$s^\vartheta \varepsilon_\vartheta(s) - \frac{\sigma_0}{E_\vartheta} = \frac{\sigma_0}{s^\vartheta \eta_\vartheta}, \quad (17)$$

which reduces to

$$\varepsilon_\vartheta(t) = \sigma_0 \left( \frac{1}{E_\vartheta} + \frac{1}{\eta_\vartheta} \frac{t^\vartheta}{\Gamma(1+\vartheta)} \right). \quad (18)$$

From Eq. (7), the fractal creep compliance of the non-differentiability type reads as follows:

$$J_\vartheta(t) = \frac{1}{E_\vartheta} + \frac{1}{\eta_\vartheta} \frac{t^\vartheta}{\Gamma(1+\vartheta)} \quad (19)$$

and its creep response is represented in Fig. 4.




Fig. 4 – The creep response of the fractal Maxwell element for  $\vartheta \ln 2 / \ln 3$ .

When the fractal body is subjected to the following condition:

$$\varepsilon_\vartheta(t) = \varepsilon_0 H(t), \quad (20)$$

where  $\varepsilon_0$  is the a constant strain, Eq. (14) can be written in the form:

$$\frac{\sigma_\vartheta(t)}{\eta_\vartheta} + \frac{1}{E_\vartheta} \frac{d^\vartheta \sigma_\vartheta(t)}{dt^\vartheta} = 0. \quad (21)$$

Taking the LFLT of both sides of Eq. (19), we have

$$\frac{\sigma_\vartheta(s)}{\eta_\vartheta} + \frac{1}{E_\vartheta} \left( s^\vartheta \sigma_\vartheta(s) - E_\vartheta \varepsilon_0 \right) = 0, \quad (22)$$

which leads us to

$$\sigma_{\vartheta}(s) = \frac{E_{\vartheta}\varepsilon_0}{s^{\vartheta} + \frac{E_{\vartheta}}{\eta_{\vartheta}}}. \quad (23)$$

For  $\sigma_0 = E_{\vartheta}\varepsilon_0$ , we find from Eq. (23) that

$$\sigma_{\vartheta}(t) = \sigma_0 M_{\vartheta} \left( -\frac{E_{\vartheta}}{\eta_{\vartheta}} t^{\vartheta} \right). \quad (24)$$

In view of Eq. (49), the fractal relaxation modulus of the non-differentiability type can be written as follows:

$$G_{\vartheta}(t) = E_{\vartheta} M_{\vartheta} \left( -\frac{t^{\vartheta}}{\tau_m} \right), \quad (25)$$

where  $\tau_m$  given by

$$\tau_m = \eta_{\vartheta} / E_{\vartheta}$$

is the relaxation time, and its relaxation response is displayed in Fig. 5.




Fig. 5 – The relaxation response of the fractal Maxwell element for  $\vartheta = \ln 2 / \ln 3$ .

## 5. FRACTAL KELVIN-VOIGT ELEMENT VIA LFD

The fractal Kelvin-Voigt element, which is used to represent the fractal solid, consists of the fractal spring and dashpot in parallel as illustrated in Fig. 6.




Fig. 6 – The representation of the fractal Kelvin-Voigt element involving LFD.

The constitutive equation of the fractal Kelvin-Voigt element within LFD takes the following form:

$$\eta_\vartheta \frac{d^\vartheta \varepsilon_\vartheta(t)}{dt^\vartheta} + E_\vartheta \varepsilon_\vartheta(t) = \sigma_\vartheta(t), \quad (26)$$

where  $\eta_\vartheta$  is the fractal viscosity,  $E_\vartheta$  is the Young's modulus,  $\varepsilon_\vartheta(t)$  is the fractal strain, and  $\sigma_\vartheta(t)$  is the fractal stress.

When the fractal body is subjected to the following condition:

$$\sigma_\vartheta(t) = \sigma_0 H(t), \quad (27)$$

where  $\sigma_0$  is the a constant stress, we deduce from Eq. (27) that

$$\eta_\vartheta \frac{d^\vartheta \varepsilon_\vartheta(t)}{dt^\vartheta} + E_\vartheta \varepsilon_\vartheta(t) = \sigma_0 H(t). \quad (28)$$

Thus, in terms of the LFLT of Eq. (27), we have

$$\eta_\vartheta \left( s^\vartheta \varepsilon_\vartheta(s) - \frac{\sigma_0}{E_\vartheta} \right) + E_\vartheta \varepsilon_\vartheta(s) = \frac{\sigma_0}{s^\vartheta}, \quad (29)$$

which implies that

$$\varepsilon_\vartheta(s) = \frac{\sigma_0}{E_\vartheta} \left( \frac{1}{s^\vartheta} - \frac{1}{s^\vartheta + \frac{E_\vartheta}{\eta_\vartheta}} \right). \quad (30)$$

From Eq. (30), the fractal strain of the solid is written in the following form:

$$\varepsilon_\vartheta(t) = \frac{\sigma_0}{E_\vartheta} \left( 1 - M_\vartheta \left( -\frac{t^\vartheta}{\tau_K} \right) \right), \quad (31)$$

where  $\tau_K = E_\vartheta/\eta_\vartheta$  is the retardation time.

In view of Eq. (30), the creep compliance of the fractal Kelvin-Voigt element can be written as follows:

$$J_\vartheta(t) = \frac{1}{E_\vartheta} \left( 1 - M_\vartheta \left( -\frac{t^\vartheta}{\tau_K} \right) \right) \quad (32)$$

and its creep response is represented in Fig. 7.




Fig. 7 – The creep response of the fractal Kelvin-Voigt element for  $\vartheta = \ln 2 / \ln 3$ .




Fig. 8 – The relaxation response of the fractal Kelvin-Voigt element for  $\vartheta = \ln 2 / \ln 3$ .

When the fractal body is subjected to the following condition:

$$\varepsilon_{\vartheta}(t) = \varepsilon_0 H(t), \quad (33)$$

where  $\varepsilon_0$  is the a constant strain, we find from Eq. (26) that

$$\sigma_{\vartheta}(t) = E_{\vartheta}\varepsilon_0 + \eta_{\vartheta}\varepsilon_0 \frac{d^{\vartheta}H(t)}{dt^{\vartheta}}, \quad (34)$$

which yields the relaxation modulus of the fractal Kelvin-Voigt element as follows:

$$G_{\vartheta}(t) = E_{\vartheta} + \eta_{\vartheta} \frac{d^{\vartheta}H(t)}{dt^{\vartheta}}, \quad (35)$$

where  $d^{\vartheta}H(t)/dt^{\vartheta}$  is the delta function defined on fractal sets [26], and its relaxation response is displayed in Fig. 8.

## 6. CONCLUSION

In our present investigation, we firstly addressed the LFRMs for the fractal Maxwell fluid and Kelvin-Voigt solid via spring and dashpot elements of non-differentiability type in the one-dimensional case based on the theory of LFC. The fractal creep compliance and relaxation modulus of the fractal Maxwell and Kelvin-Voigt elements were graphically presented with the aid of the LFLT. The viscoelastic behaviors of non-differentiability type in the fractal material were also discussed. For  $\vartheta = 1$ , the proposed models turn out to be those of the classical rheological behaviors (see [36] for further details of these ideas and concepts). Our presentation here may open up a new perspective in description of the fractal rheological behaviors of the real materials.

*Acknowledgements.* This work was supported by the State Key Research Development Program of the People's Republic of China (Grant No. 2016YFC0600705) and the Priority Academic Program Development of the Jiangsu Higher Education Institutions (PAPD2014).

## REFERENCES

1. F. Mainardi, *Fractional Calculus and Waves in Linear Viscoelasticity: An Introduction to Mathematical Models*, World Scientific, Hong Kong, HK, 2010.
2. G. S. Blair, *The role of psychophysics in rheology*, Journal of Colloid Science, **2**, 1, 21–32 (1947).
3. A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and Applications of Fractal Differential Equations*, North-Holland Mathematical Studies, Vol. 204, Elsevier (North-Holland) Science Publishers, Amsterdam, 2006.
4. A. S. Balankin, *A continuum framework for mechanics of fractal materials I: from fractional space to continuum with fractal metric*, The European Physical Journal B, **88**, 4, 90 (2015).
5. A. S. Balankin, *A continuum framework for mechanics of fractal materials II: elastic stress fields ahead of cracks in a fractal medium*, The European Physical Journal B, **88**, 4, 91 (2015).
6. V. E. Tarasov, *Three-dimensional lattice models with long-range interactions of Grünwald–Letnikov type for fractional generalization of gradient elasticity*, Meccanica, **51**, 1, 125–138 (2016).

7. V. E. Tarasov, *Discrete model of dislocations in fractional nonlocal elasticity*, Journal of King Saud University-Science, **28**, 1, 33–36 (2016).
8. M. A. Abdelkawy, M. A. Zaky, A. H. Bhrawy, D. Baleanu, *Numerical simulation of time variable fractional order mobile-immobile advection-dispersion model*, Romanian Reports in Physics, **67**, 773–791 (2015).
9. A. H. Bhrawy, *A new spectral algorithm for time-space fractional partial differential equations with subdiffusion and superdiffusion*, Proceedings of the Romanian Academy Series A, **17**, 39–47 (2016).
10. K. Nouri, S. Elahi-Mehr, L. Torkzadeh, *Investigation of the behavior of the fractional Bagley-Torvik and Basset equations via numerical inverse Laplace transform*, Romanian Reports in Physics, **68**, 503–514 (2016).
11. W. M. Abd-Elhameed, Y. H. Youssri, *Spectral solutions for fractional differential equations via a novel Lucas operational matrix of fractional derivatives*, Romanian Journal of Physics, **61**, 795–813 (2016).
12. A. Agila, D. Baleanu, R. Eid, B. Irfanoglu, *Applications of the extended fractional Euler-Lagrange equations model to freely oscillating dynamical systems*, Romanian Journal of Physics, **61**, 350–359 (2016).
13. A. N. Gerasimov, *A generalization of linear laws of deformation and its application to problems of internal friction*, Akad. Nauk SSSR Prikl. Mat. Meh., **12**, 251–260 (1948) (in Russian).
14. M. Caputo, F. Mainardi, *Linear models of dissipation in anelastic solids*, Rivista del Nuovo Cimento (Ser. II), **1**, 161–198 (1971).
15. W. Smit, H. De Vries, *Rheological models containing fractional derivatives*, Rheologica Acta, **9**, 4, 525–534 (1970).
16. F. Gao, X.-J. Yang, *Fractional Maxwell fluid with fractional derivative without singular kernel*, Thermal Science, **20**, 3, s871–s877 (2016).
17. F. Mainardi, R. Gorenflo, *Time-fractional derivatives in relaxation processes: a tutorial survey*, Fractional Calculus and Applied Analysis, **10**, 1, 269–308 (2007).
18. L. B. Eldred, W. P. Baker, A. N. Palazotto, *Kelvin-Voigt versus fractional derivative model as constitutive relations for viscoelastic materials*, Aiaa Journal, **33**, 3, 547–550 (1995).
19. R. Metzler, T. F. Nonnenmacher, *Fractional relaxation processes and fractional rheological models for the description of a class of viscoelastic materials*, International Journal of Plasticity, **19**, 7, 941–959 (2003).
20. R. L. Bagley, P. J. Torvik, *On the fractional calculus model of viscoelastic behavior*, Journal of Rheology, **30**, 1, 133–155 (1986).
21. F. Mainardi, *An historical perspective on fractional calculus in linear viscoelasticity*, Fractional Calculus and Applied Analysis, **15**, 4, 712–717 (2012).
22. Z. J. Meng, Y. M. Zhou, H. Q. Wang, *Local fractional variational iteration algorithm iii for the diffusion model associated with non-differentiable heat transfer*, Thermal Science, **20**, 3, S781–S784 (2016).
23. Z. J. Meng, Y. M. Zhou, D. M. Mei, *On three-dimensional incompressible Navier-Stokes fluid on cantor sets in spherical Cantor type coordinate system*, Thermal Science, **20**, 3, S853–S858 (2016).
24. X.-J. Yang, J. A. T. Machado, D. Baleanu, C. Cattani, *On exact traveling-wave solutions for local fractional Korteweg-de Vries equation*, Chaos: An Interdisciplinary Journal of Nonlinear Science, **26**, 8, 110–118 (2016).
25. X.-J. Yang, H. M. Srivastava, *An asymptotic perturbation solution for a linear oscillator of free damped vibrations in fractal medium described by local fractional derivatives*, Communications

- in *Nonlinear Science and Numerical Simulation*, **29**, 1-3, 499–504 (2015).
26. X.-J. Yang, D. Baleanu, H. M. Srivastava, *Local Fractional Integral Transforms and Their Applications*, Academic Press (Elsevier Science Publishers), Amsterdam, Heidelberg, London and New York, 2016.
  27. M. Sarikaya, H. Budak, *Generalized Ostrowski type inequalities for local fractional integrals*, *Proceedings of the American Mathematical Society*, **145**, 4, 1527–1538 (2016).
  28. K. M. Kolwankar, A. D. Gangal, *Local fractional Fokker-Planck equation*, *Physical Review Letters*, **80**, 2, 214–217 (1998).
  29. K. M. Kolwankar, A. D. Gangal, *Hölder exponents of irregular signals and local fractional derivatives*, *Pramana*, **48**, 1, 49–68 (1997).
  30. A. Carpinteri, P. Cornetti, K. M. Kolwankar, *Calculation of the tensile and flexural strength of disordered materials using fractional calculus*, *Chaos Solitons and Fractals*, **21**, 3, 623–632 (2004).
  31. C. Cattani, H. M. Srivastava, X.-J. Yang, *Fractional Dynamics*, Emerging Science Publishers (De Gruyter Open), Berlin and Warsaw, 2015.
  32. D. Kumar, J. Singh, D. Baleanu, *A hybrid computational approach for KleinCGordon equations on Cantor sets*, *Nonlinear Dynamics*, **87**, 1, 511–517 (2017).
  33. H. Jafari, H. K. Jassim, F. Tchier, D. Baleanu, *On the approximate solutions of local fractional differential equations with local fractional operators*, *Entropy*, **18**, 4, 1–12 (2016).
  34. J.-D. Chen, H.-P. Li, *The non-differentiable solution for local fractional Laplace equation in steady heat-conduction problem*, *Thermal Science*, **20**, 3, S769–S772 (2016).
  35. Y. Zhang, D. Baleanu, Xiao-Jun Yang, *New solutions of the transport equations in porous media within local fractional derivative*, *Proceedings of the Romanian Academy Series A*, **17**, 230–236 (2016).
  36. G. T. Mase, R. E. Smelser, G. E. Mase, *Continuum Mechanics for Engineers*, Third edition, Series on Applied Mathematics and Computational Mechanics, CRC Press (Taylor and Francis Group), Baton Roca, 2009.